

Aerobic Vinyl Chloride Biodegradation to Support Anaerobic Chloroethene Reduction

Presented By Tony Danko, Ph.D.

Naval Facilities Engineering Command (NAVFAC)
Engineering and Expeditionary Warfare Center (EXWC)


Objective



 Present and overview of aerobic VC biodegradation its potential impact on the attenuation of VC and other cVOCs


Conventional CSM for Chloroethenes





Anaerobic Reductive Dechlorination of Chloroethenes





Dehalobacter
Desulfitobacterium
Desulfuromonas

Dehalococcoides

Dehalogenimonas Etc. <u>Dehalococcoides</u> <u>Dehalogenimonas</u>


Microorganisms

Source: Frank Löffler

Aerobic Dechlorination Processes



Genes




EtnC = Alkene Monooxygenase EtnE = Epoxyalkane CoM Transferase

Aerobic Dechlorination Processes



Genes




Etheneotrophs VC-assimilators

EtnC = Alkene Monooxygenase EtnE = Epoxyalkane CoM Transferase


Metabolism versus Cometabolism





Aerobic Dechlorination Processes





Propanotrophs Methanotrophs etc.

TOD = Toluene Dioxygenase sMMO/pMMO = soluble/particulate Methane Monooxygenase PHE = Phenol Hydroxylase RMO = Toluene Monooxygenase RDEG 2 = Toluene Monooxygenase

EtnC = Alkene Monooxygenase

Microorganisms


Anaerobic and Aerobic Degraders



 Previous work at 5 different cVOC sites with different concentrations and biodegradation rates but generally low in DO and ORP

Liang et al., 2017

- Found high concentrations of aerobic VC genes/transcripts (etnC/etnE) + anaerobic VC genes/transcripts (vcrA/bvcA) associated with VC
- ONot true with methanotrophs




NOTE: K_s for O_2 for VC degraders is ~ 0.16 mg/L!

Coleman et al., 2002

Anaerobic and Aerobic Degraders



cVOC site



Parameter	Avg. sample (39) concentrations
PCE + TCE	292 μg/L
cDCE	2651 μg/L
VC	1938 μg/L
Ethene	218 μg/L
Methane	2621 μg/L
DO	0.74 mg/L
ORP	15 mV


Cryocore sampling






Gas Concentrations





Anaerobic and Aerobic Degraders





Knowledge Check



 Biotransformation of VC only occurs via anaerobic processes (Dehalococcoides)

TRUE

FALSE

• A variety of primary substrates can be present at a site that can drive cometabolism

TRUE

FALSE

Summary



- Both anaerobic and aerobic processes are able to degrade chloroethenes at sites
- Aerobic microorganisms are active in anaerobic zones
- Both metabolic and cometabolic processes may be important
- Biomarkers are available to measure the potential for anaerobic and aerobic activity
- Data should be integrated into existing site data

Contacts and Questions



Points of Contact

NAVFAC LIST FEC: Tony Danko, NAVFAC EXWC

anthony.danko@navy.mil

NAVFAC LIST FEC: Arun Gavaskar, NAVFAC EXWC

arun.gavaskar@navy.mil

Questions?

Supplemental Information



Advances in Microbial Characterization for MNA & Bioremediation (Open Environmental Restoration Resources (OER2) Webinar):

https://www.navfac.navy.mil/navfac worldwide/specialty centers/exwc/products a

nd_services/ev/erb/oer2.html



Relationships between the Abundance and Expression of Functional Genes from Vinyl Chloride (VC)-Degrading Bacteria and Geochemical Parameters at VC-Contaminated Sites

Y. Liang, X. Liu, M. Singletary, K. Wang, and T. E. Mattes https://pubs.acs.org/doi/10.1021/acs.est.7b03521

NAVFAC Environmental Restoration and BRAC Website

→ Chlorinated Solvents

https://www.navfac.navy.mil/navfac_worldwide/specialty_centers/exwc/products_a nd_services/ev/erb/chlor-solv.html#resources